

RESPONSE OF DATE PALM (*Phoenix dactylifera* L.) OFF SHOOTS TO BIOSTIMULANT EM AND CHEMICAL FERTILIZER UNDER NURSERY CONDITION

(Received: 14.9.2015)

By
L. M. Abdel-Galeil

*Central Laboratory of Date palm Research and Development,
Agricultural Research Center, Giza, Egypt*

ABSTRACT

A trial was carried out at the Experimental Farm of the Hort. Res. Inst., A.R.C., Giza, Egypt during 2014 and 2015 seasons to study the effect of individual application of Em biostimulant solution at 25 ml/l, NPK complete chemical fertilizer (20 : 20 : 20 + microelements) at the rates of 0, 2.5, 5.0, 7.5 and 10.0 g/bag and their combinations on growth and chemical composition of 2 year – old off shoots of date palm (*Phoenix dactylifera* L.) cv. Siwi planted in 50-cm-diameter plastic bags filled with about 35 kg of a mixture of sand and clay (2 : 1, v/v) under the full sun.

The results showed that all fertilization treatments used markedly improved all vegetative growth characteristics with various significant levels when compared to the control in both seasons. The results indicated also that EM solution alone gave means closely near those gained by NPK complete fertilizer at 5 g/bag with few exceptions in the two seasons. Moreover, increasing the application rate of NPK fertilizer to 7.5 g/bag or more did not cause a pronounced increment in growth, but combining between EM treatment and NPK fertilizer at any rate did so, with the superiority of combining between 25 ml/l EM and 5g NPK/bag, as this combination gave the highest means over other combined treatments in both seasons. Similar observations were also attained concerning the leaflet content of N, P, K, total indoles, chlorophylls a and b and carotenoids.

Hence, it can be recommended to treat the 2year old offshoots of date palm cv. Siwi cultivated in 50cm diameter plastic bags with EM solution at 25ml/l and NPK complete fertilizer at 5 g/bag to improve growth and quality of such offshoots before transferring to the permanent field.

Key words: Date Palm-*Phoenix dactylifera* L., EM biostimulant, NPK fertilization.

1. INTRODUCTION

Phoenix dactylifera L., date palm (Fam. Palmaceae) is one of the common and popular fruit grown in tropical and subtropical areas. There are many cultivars of date palm, some of them give semi-dry fruits, among which cv. Siwi. It is one of the most famous and distributed cultivars in Egypt, especially at the governorates of New Valley, Wahat, Giza and Fayoum. Each cv. Siwi date palm tree yields about 90 up to more than 150 kg fruits per year. The fruits are somewhat big-sized, reaching 3.5-4.0cm in length and 2.0-2.5cm in diameter. They are yellow at mature stage as they can be consumable at this phase. After drying, these fruits turn to deep brown. It is among the highly qualified cultivars for processing and packing as pressed dates. The fruits can be stored for long time after harvest.

It is well known that all palm plants grow

slowly, and fertilizing them with either chemical or bio-preparations usually enhance growth, especially at the early stages. In this concern, Abdel-Galeil *et al.* (2010) found that the survival and rooting percentages, root length and No. roots/ offshoot of date palm cv. Zaghloul were linearly increased with increasing the level of commercial liquid fertilizer (10N: 10P: 10K + micro-elements) from 5 to 10 ml/l. A similar trend was also attained regarding leaf length, No. new formed leaves/ offshoot, fresh and dry weights of the new formed leaves, as well as the percentages of N, P and K in the new leaves. Likewise, the growth parameters were linearly increased with increasing the level of commercial liquid fertilizer (10N : 10P : 10K + microelements) from 5 to 10 ml/l. A similar trend was also attained regarding leaf length, No. new formed leaves / offshoots, fresh and dry weights of the new formed leaves, as well as the

percentages of N, P and K in the new leaves. However, Abdel-Galeil (2010a) on date palm cv. Malacabe mentioned that combining NPK liquid fertilizer (10 : 10 : 10+ microelements) at 2 ml/l as a foliar spray and at 20 ml/l as a soil drench greatly improved trunk length, No. leaves/plantlet and leaf width. Connecting between potassien-N (K-N) at 2 ml/l as a foliar spray and NPK liquid fertilizer (10 : 10 : 10+ microelements) at 20 ml/l as a soil drench significantly increased trunk length, No. leaves/plant, leaf width, as well as content of chlorophyll a and b, carotenoids, N, P and K in the leaves of cv. Sakkoty plant (Abdel-Galeil, 2010a).

On ornamental plants, Agina *et al.* (2005) on *Ficus macrocarpa* cv. Hawaii, reported that foliar application of kristalon (19 : 19: 19+ micronutrients) markedly improved vegetative growth, fresh and dry weights of leaves, stem and roots, as well as pigments and minerals content in the leaves. Similarly, the results were recorded by Kandeel *et al.* (2002) on *Melia azedarach*, Sarhan *et al.* (2002) on *Taxodium disticum*, Gad (2003) on *Ficus benjamina*, El-Sayed *et al.* (2008) on *Ficus macrocarpa* cv. Hawaii, Abdel-Fattah *et al.* (2009) on *Dracaena* and *Ruscus*, Shahin *et al.* (2012) on *Schefflera* and *Euonymus*, El-Fouly *et al.* (2014) on *Ficus deltoidea* and Shahin *et al.* (2014) on *Merremia dissecta*.

A commercial Japanese product, EM is a biostimulant that contains more than 60 selected strains of "Effective Microorganisms", viz., Photosynthetic bacteria, lactic acid bacteria, yeast, actinomycetes and various fungi that improve growth and health of plants (Primavesi, 1999). Janas (2009) revealed that effective microorganisms (EM) is characterized by a wide spectrum of activity and complex effect on plant living environment. Thus, it may be used as foliar treatments, on the seeds or in soil application. Its effects on inducing plant disease resistance, yield creating and protective were observed in many industrial, medicinal and ornamental plant species. It also creates humus substance and regulates basic relations in the soils. Therefore, the EM bispreparation is used

in many countries, on a large scale, in organic production of agricultural crops. In this regard, Thach *et al.* (1999) indicated that treatment orchids with EM led to larger stems, darker green leaves and accelerating flowering in *Dendrobium* plants. On jojoba, Sarhan *et al.* (2007) noticed that Biomagic (a commercial biopromotor) at 10g/l and inoculation with a mixture of *Azotobacter* and *Bacillus* significantly increased vegetative growth parameters, pigments content, total carbohydrates, N, P, K, Fe, Zn and Cu in the leaves. On the same line, were those results recorded by El-Seginy (2006) on pear and apricot and El-Sayed (2012) on paspalum turf.

This work was set out to discover the beneficial effects of both chemical fertilizer and biostimulant, alone or in combination on growth and quality of offshoots of date palm cv. Siwi.

2. MATERIALS AND METHODS

This study was conducted at the Experimental Farm of the Hort Res. Inst., ARC, Giza, Egypt during two consecutive seasons of 2014 and 2015 to reveal the effect of complete chemical fertilizer and EM biostimulant, solely or in combination treatments on growth and chemical composition of Siwi cultivar offshoots date palm.

Two-year-old uniform offshoots were selected carrying about 4-5 leaves. The bases of the selected offshoots were first dipped in a 0.5% solution of Topsin-M, 70% WP (Sumitomo Chemical Co., Ltd., Osaka, Japan) for 30 minutes, and then planted on April, the 1st for the two studied seasons in 50-cm-diameter plastic bags (one offshoot/bag) filled with about 35 kg of sand and clay soil mixture (2:1, by volume). Some physical and chemical properties of the sand and clay used in both seasons were determined according to the standard methods described by Richards (1954) and illustrated in Table (1).

After one week from planting, the planted offshoots were irrigated with 10 liters of fresh water/bag. On the 15th of April, the offshoots received the following treatments:

Table (1): Some physical and chemical properties of the used sand and clay in both seasons.

Soil type	Particle size distribution (%)				S.P	E.C. (ds/m)	pH	Cations (meq/l)				Anions (meq/l)		
	Coarse sand	Fine sand	Silt	Clay				Ca ⁺⁺	Mg ⁺⁺	Na ⁺	K ⁺	HCO ₃ ⁻	Cl ⁻	SO ₄ ⁻
Sand	18.72	71.28	4.76	5.34	21.83	1.58	8.20	2.65	2.48	21.87	0.78	3.85	13.00	10.93
Clay	7.46	16.75	34.53	40.89	41.76	2.18	8.33	16.93	9.33	20.44	0.37	3.82	1.46	41.79

1. No fertilization, referred to as the control.
2. EM biostimulant solution consisting of 0.5l EM+ 1l molasses +20l well water (25 ml EM/l) was fermented for one week under an aerobic conditions before application as a soil drench and 500 ml of EM suspension solution were added to a bag.
3. A commercial complete chemical fertilizer 20N: 20P: 20K+ microelements) known as Ectaful was added also as a soil application at the rates of 2.5, 5.0, 7.5 and 10.0 g/bag. The chemical composition of such fertilizer is shown in Table (2).
4. EM solution at 25 ml/l was combined with each level of NPK commercial fertilizer to form 4 combinations as follows:
 - a. EM at 25ml/l + NPK fertilizer at 2.5g/ bag.
 - b. EM at 25ml/l + NPK fertilizer at 5.0g/ bag.
 - c. EM at 25ml/l + NPK fertilizer at 7.5g/ bag.
 - d. EM at 25ml/l + NPK fertilizer at 10.0g/ bag.

Data were then tabulated, and the morphological ones were subjected to analysis of variance according to SAS Institute Program (1994) was used for statistical analysis, whereas Duncan's Multiple Range Test (1955) was employed to verify the differences among the means of different treatments.

3. RESULTS AND DISCUSSION

3.1. Effect of fertilization treatments on

3.1.1. Vegetative growth parameters

It is evident from Table (3) that all fertilization treatments employed in general improved, all vegetative growth parameters with various significant differences when compared to the untreated control in both seasons. It was also noticed that EM alone gave means closely near to those of NPK fertilizer at 5 g/bag, with few exceptions in the two seasons, while combining between these two treatments gave the utmost high means over the control and all

Table (2): The chemical composition of the commercial NPK fertilizer used in the two seasons.

Component	Value	Component	Value	Component	Value
Nitrogen (N)	20%	Chelated Fe	700ppm	Chelated Zn	140ppm
Phosphorus (P)	20%	Chelated Mn	420ppm	Chelated Mo	140ppm
Potassium (K)	20%	Chelated Cu	160ppm	Boron (B)	220ppm

The previous treatments were applied 4 times commencing from mid April till mid October (one every two months). The experimental treatments in both seasons were arranged in a complete randomized design (Mead *et al.*, 1993) and the treatments were replicated 3 times with 3 offshoots for each replicate, *i.e.* 10 treatments X3 replicates X3 offshoots = 90 offshoots for each season. All the offshoots received the usual agricultural practices recommended for such plantation whenever needed.

At the end of each season (the end of October), the following measurements were recorded: length of the first new formed leaf (cm), number of the new formed leaves / offshoot, trunk length and circumference (cm) and fresh and dry weights of the first new leaf (g). In fresh leaf samples taken from the leaflets of the first new leaf, photosynthetic pigments (chlorophyll a, b and carotenoids, mg/g f.w.) and the total indoles were determined according to the methods of Moran (1982) and A.O.A.C. (1980), respectively, while in dry leaf samples, the percentages of nitrogen (using micro-Kjeldahl method described by Pregli, 1945) and potassium (using flame photometer set as explained by Jackson, 1973) were evaluated.

other individual or combined treatments in both seasons. Increasing the rate of NPK fertilizer to 7.5g /bag or more did not induce valuable additional increment in the growth, whereas combining between EM treatment and NPK fertilizer at any level did so, especially between 25 ml/l EM and 5 g NPK/bag combination, which was superior over all other sole and combined treatments in both seasons, as mentioned before.

Thus, using NPK complete fertilizer with EM biostimulant seemed to be valuable for enhancing growth through providing the plants with macro-and micro-nutrients necessary for good and healthy growth, inducing plant disease resistance by EM, which also creates humus substances and regulates basic relations in the soil (Janas, 2009). Moreover, EM may play a role in enhancing the enzymatic systems in the plant tissues, and consequently activating growth (Thach *et al.*, 1999).

Means within columns having the same letters are not significantly different according to Duncan multiple range test (1955)

The present results are in harmony with those postulated by Abdel-Galeil (2010a and b) on date palm cvs. Malacabe and Sakkoty,

Table (3): Effect of fertilization treatments on some vegetative growth traits of *Phoenix dactylifera l.* cv . Siwi during 2014 and 2015 seasons.

Treatments	Length of the first new leaf (cm)	No. the new leaves/ off shoot	Trunk length (cm)	Trunk circumference at the base (cm)	F.W. the first new leaf (g)	D.W. the first new leaf (g)
First seasons 2014						
Control	63.00 e	2.00 c	28.67 f	25.37 f	64.31 d	30.50 c
EM at 25ml/l (A)	75.40 cd	3.00 b	36.00 d	31.33 e	68.40 c	33.16 bc
NPK at 2.5g/bag (B)	72.33 d	3.00 b	31.50 e	30.50 e	66.35 cd	32.18 cb
NPK at 5.0g/bag (C)	75.65 cd	3.33 b	38.46 cd	31.60 e	68.50 c	33.44 bc
NPK at 7.5g/bag (D)	78.00 c	3.33 b	45.00 bc	39.73 c	73.33 b	35.20 b
NPK at 10.0g/bag (E)	78.33 c	3.00 b	46.10 b	37.00 d	72.38 b	35.00 b
A + B	76.10 cd	3.33 b	45.33 b	44.00 b	70.45 bc	34.56 bc
A + C	113.33 a	4.00 a	53.67 a	51.00 a	80.34 a	41.78 a
A + D	92.48 b	3.33 b	38.70 cd	42.71 bc	77.50 ab	36.85 b
A + E	93.21 b	3.33 b	41.33 c	39.80 c	73.44 b	35.22 b
Second seasons 2015						
Control	56.33 g	2.33 c	31.30 e	26.33 e	60.33 e	30.00 d
EM at 25ml/l (A)	68.10 e	3.00 bc	40.00 c	33.50 d	68.14 de	31.34 cd
NPK at 2.5g/bag (B)	61.79 f	3.00 bc	35.67 d	31.67 d	67.13 ed	33.18 c
NPK at 5.0g/bag (C)	71.50 ed	3.33 b	43.33 b	42.33 b	70.45 d	35.23 bc
NPK at 7.5g/bag (D)	74.00 d	3.00 bc	37.90 dc	41.00 b	73.40 c	36.20 b
NPK at 10.0g/bag (E)	72.43 de	3.00 bc	36.50 d	37.38 c	72.12 cd	35.03 bc
A + B	76.00 cd	3.33 b	41.86 bc	36.51 c	75.33 cb	35.11 bc
A + C	95.97 a	4.33 a	50.98 a	53.00 a	85.48 a	42.16 a
A + D	88.33 b	3.00 bc	40.36 c	41.83 b	78.43 b	37.10 b
A + E	79.00 c	3.00 bc	38.39 cd	40.33 b	73.12 c	35.14 bc

Means within colams having be letters are not significantly difficent according to Duncan multiple rang test

Kandeel *et al.* (2002) on *Melia azedarach*, Agina *et al.* (2005) on *Ficus macrocarpa* cv. Hawaii, Abdel-Fattah *et al.* (2009) on *Dracaena* and *Ruscus* and El-Sayed (2012) who elicited that complete fertilizer (19 : 19 : 19+ micronutrients) at 2g/pot plus EM at 1ml/l as a soil drench gave the best growth, density and color in seashore paspalum turf.

3.1.2. Leaf chemical composition

The data averaged in Table (4) show that the percentages of N, P and K were markedly increased in the leaves of the plants treated with either NPK complete fertilizer or EM biostimulant, alone or in combination over the percentages gained by the untreated plants in the two seasons. The least records of these nutrients in both seasons were found due to EM solution (25 ml/l) and the low rates of NPK fertilizer when each of them was applied alone. However, combining between EM and the complete fertilizer gave higher records, especially when the combining between EM at 25 ml/l and NPK fertilizer at 5 g/bag, as this combination registered the highest content at all in both

seasons. A similar response occurred as well in respect of the total indole content (ppm) and pigments content (mg/g F.W.) in the two seasons, except for EM solution and NPK fertilizer at 10.0g/ bag treatments which gave less content of chlorophyll b in the first season than the control (0.055 and 0.058mg/g f .w., respectively against 0.064mg/g F.W. for control), and EM solution alone that recorded less content of carotenoids than control in the two seasons (0.058 mg/g f .w. against 0.070 mg/g f .w. for control in the 1st season, and 0.061 mg/g f.w. against 0.072mg/g f.w. for the control in the 2nd one).

In general, the prevalence in all previous constituents' content was for the combined treatment of 25 ml/l EM+ 5g/bag NPK complete fertilizer which scored the utmost high records in both seasons.

These findings are reasonable because the presence of EM, as a biostimulant reinforce the beneficial effects of NPK fertilizer which supply the plants with the different nutrients necessary for good growth, besides the role of

Table (4): Effect of fertilization treatments on the chemical composition of *Phoenix dactylifera* L.cv. Siwi leaves during 2014 and 2015 seasons.

Treatments	N %	P %	K %	Total indoles (ppm)	Pigments content (mg/ g f.w.)		
					Chlo.a	Chlo.b	Carotenoids
First seasons 2014							
Control	1.78	0.11	1.07	0.139	0.099	0.064	0.070
EM at 25ml/l (A)	2.47	0.15	1.19	0.226	0.182	0.055	0.058
NPK at 2.5g/bag (B)	2.33	0.16	1.26	0.198	0.120	0.068	0.073
NPK at 5.0g/bag (C)	2.56	0.28	1.50	0.377	0.156	0.068	0.097
NPK at 7.5g/bag (D)	2.90	0.30	1.50	0.240	0.133	0.079	0.099
NPK at 10.0g/bag (E)	2.71	0.26	1.63	0.293	0.137	0.058	0.096
A + B	2.50	0.31	1.44	0.232	0.376	0.087	0.100
A + C	3.31	0.43	1.72	0.521	1.182	0.183	0.127
A + D	2.95	0.30	1.58	0.395	0.490	0.076	0.093
A + E	2.95	0.24	1.50	0.253	0.286	0.065	0.089
Second seasons 2015							
Control	1.81	0.10	1.13	0.143	0.095	0.062	0.072
EM at 25ml/l (A)	2.50	0.13	1.20	0.231	0.174	0.074	0.061
NPK at 2.5g/bag (B)	2.37	0.15	1.27	0.198	0.124	0.064	0.077
NPK at 5.0g/bag (C)	2.61	0.21	1.45	0.373	0.231	0.071	0.090
NPK at 7.5g/bag (D)	2.95	0.27	1.51	0.393	0.190	0.076	0.091
NPK at 10.0g/bag (E)	2.70	0.30	1.60	0.370	0.135	0.084	0.086
A + B	2.63	0.30	1.50	0.391	0.277	0.090	0.091
A + C	3.18	0.36	1.76	0.458	1.310	0.181	0.115
A + D	2.87	0.28	1.63	0.239	0.383	0.103	0.101
A + E	2.90	0.25	1.51	0.352	0.381	0.092	0.094

EM in increasing the surface unit area of root length and hence enhancing the root hair branching with an eventual increase in acquisition of nutrients from the soil solution (Primavesi,1999). These positive responses were also acknowledged by numerous investigators such as Abdel-Galeil *et al.* (2010) on date palm cv. Zaghoul, Gad (2003) on *Ficus benjamina*, El-Sayed *et al.* (2008) on *Ficus macrocarpa* cv. Hawaii and Shahin *et al.* (2014) on *Merremia dissecta*. In this regard, El-Seginy (2006) declared that the use of organic fertilizer (10 : 10 : 10+ micro-nutrients and humates) and EM biostimulant on young Le Conte pear and Canino apricot trees grown in calcareous soil gave vigorous growth and increased leaf nutrients content. It also decreased the cost production and boosted the income.

From the aforementioned results, it can be advised to treat the 2-year-old offshoots of date palm cv. Siwi with EM solution (25ml/l) plus NPK complete fertilizer (5 g/bag) to improve their vegetative growth and quality.

4. REFERENCES

- Abdel-Fattah G. H., El-Sayed B. A. and Khenizy S. A. M. (2009). Response of *Dracaena* and *Ruscus* plants to humic acid and biofertilizer supply. Ann. Agric. Sci., Moshtohor, 47(1): 111-119.
- Abdel-Galeil L. M. (2010a). Response of date palm plantlets cv. Malacabe to some fertilization treatments. J. Biol. Chem. & Environ. Sci., 5(1): 43-55.
- Abdel-Galeil L. M. (2010b). Improving the growth of date palm cv. Sakkoty, plantlets by some fertilization treatments. J. Biol. Chem. & Environ. Sci., 5(1): 109-122.
- Abdel-Galeil L. M., El-Sayed B. A. and Shahin S. M. (2010). Response of date palm cv. Zaghoul off shoots to phloroglucinol and humic acid under nursery conditions. J. Biol. Chem. & Environ. Sci., 5(4): 39-52.
- Agina Eman A. M., Shalaby H. S., El-Khayat A. S. and Korkor H. M. (2005). Effect of foliar fertilization and some growth regulators on growth and chemical composition of some ornamental plants. Proc 6th Arab Conf. Hort., 20-22 March, Dept. Hort., Fac. Agric., Ismailia, Egypt, Abst.: 10.
- A.O.A.C. (1980). Official Methods of Analysis

- of the Association Official Agricultural Chemists. 15th Ed., Arlington, Virginia USA. 22201:877-878.
- Duncan D. B. (1955). Multiple range and multiple F-tests. J. Biometrics, 11:1-42
- El-Fouly A. S., Abdel-Moniem A. M. and Ibrahim H. E. (2014). Response of the slow-growing Mistletoe Fig (*Ficus deltoidea* Jack) plant to fertilization treatments and growth activator. II. Humic acid liquid fertilizer treatment. Scientific J. Flowers & Ornam. Plants, 1(1): 25-34.
- El-Sayed B. A. (2012). Response of seashore paspalun to treatment with kristalon and biostimulant EM. Minufiya J. Agric. Res., 37(4): 935-941.
- El-Sayed B. A., El-Fouly A. S. and El-Feky A. H. (2008). Response of *Ficus macrocarpa* L. var. Hawaii transplants to some fertilization treatments. Egypt. J. Appl. Sci., 23(1): 224-231.
- El-Seginy A. M. (2006). Effect of the organic fertilizer (Actosol) and EM biostimulant on vegetative growth and chemical composition of young pear and apricot trees grown in calcareous soil. J. Agric. Sci., Mansoura Univ., 31(5): 3147-3158.
- Gad M. M. (2003). Evaluation of various potting media and fertilizer levels for commercial nursery production of *Ficus benjamina* L. Assuit J. Agric. Sci., Fac. Agric., Assuit Univ., Egypt, 34: 123-151.
- Jackson M. (1973). Soil chemical analysis. Prentice-Hall of India private Ltd. M-97, New Delhi, India, 498pp.
- Janas R. (2009). Possibilities of using effective microorganisms in organic production systems of cultivated crops. Proplemy Inzynierri Rolniczei, 17(3): 111-119.
- Kandeel Y. M., El-Tarawy M. A. and El-Mahrouk E. M. (2002). Response of *Melia azedarach* L. to fertilization and irrigation treatments in Nuberia. 2nd Inter. Conf. Hort Sci., 10-12 Sept., Kafr El-Sheikh, Tanta Univ., Egypt: 484-499.
- Mead R., Curnow R. N. and Harted A. M. (1993). Statistical methods in agriculture and experimental biology, 2nd Ed., Chapman & Hall Ltd., London 335pp.
- Moran R. (1982). Formula for determination of chlorophyllous pigments extracted with N-N-dimethyl formamide. Plant physiol., 69:1376-81
- Pregli F. (1945). Quantitative Organic Micro-Analysis, 4th Ed., J. and A. Churchill Ltd., London, P.203-209.
- Primavesi A. M. (1999). Determination of plant health by their magnetic emanation and its improvement with EM. 5th Inter. Conf. on Kyusei nature Farming, Bangkok, Thailand, 23-26 Oct., 219-225.
- Richards L. A. (1954). Diagnosis and Improvement saline and Alkali Soils. USDA, Handbook No.60.
- Sarhan A., Abou-Dahab T., Abdel-Dayem A. M. and Rabie A. R. (2002). NPK fertilization of *Taxodium disticum* seedlings. 2nd Inter conf. Hort Sci., 10-12 Sept. Kafr El-Sheikh, Tanta Univ., Egypt: 1342-1358.
- Sarhan A., El-Maadawy E. I. and Surour S.S. (2007). Effect of biopromoters and biofertilizers application on growth and chemical composition of jojoba plants. J. Agric. Sci., Mansoura Univ., Egypt, 32(8): 6575-6598.
- SAS Institute (1994). SAS/STAT User's Guides Statistics. Vers. 6.04, 4th Ed., SAS Institute Inc., Cary, N.C., USA.
- Shahin S.M., El-Tayeb H. F. and El-Sayed B. A. (2012). Effect of some media and fertilization treatments on growth and quality of some foliage pot-plants. J. Biol. Chem. & Environ. Sci., 7(1): 93-112.
- Shahin S.M., El-Sayed B. A. and El-Tayeb H. F. (2014). Improving growth and quality of *Merremia dissecta* (Jacq.) H. G. Hallier twiner by some fertilization treatments. Alex. J. Agric. Sci., 59(1): 43-49.
- Thach N. Q., Long C.A., Liet V., Trung N. V., Thanh N. X., Dichi T.V., Duong N., Tuan N. K., Xuan L.T.H. and Dar P.V. (1999). Preliminary results of EM application in Vietnam. 5th Inter. Conf. on Kyusei Nature Farming, Bangkok, Thailand, 23-26 Oct. P.254-260.

استجابة فساتل نخيل البلح للمنشط الحيوي EM والتسميد الكيماوى تحت ظروف المشتل

لبنى محمد عبد الجليل

المعمل المركزي لأبحاث وتطوير نخيل البلح - مركز البحوث الزراعية - الجيزة - مصر.

ملخص

أجريت هذه الدراسة بالمزرعة التجريبية لمعهد بحوث البساتين، مركز البحوث الزراعية، الجيزة، مصر خلال موسمي 2014، 2015 لدراسة تأثير الإضافة الفردية الأرضية لمحلول المنشط الحيوي (EM) بمعدل 25 مللى/لتر، السماد الكيماوي المتكامل NPK (20 : 20 : 20 + عناصر صغرى) بمعدل: صفر، 2.5، 5، 7.5، 10 جم/ كيس (أي لكل فسيلة) والتوليفات بينهما على النمو والتركيب الكيميائي لفساتل عمر سنتين من نخيل البلح صنف سيوي (نصف جاف) المنزرعة في أكياس بلاستيك قطرها 50 سم ملأت بحوالي 35 كجم من مخلوط الرمل والطين (بنسبة 2 : 1 حجمًا) تحت ظروف الحقل المفتوح. أوضحت النتائج أن جميع المعاملات المطبقة بهذه الدراسة أحدثت تحسناً ملحوظاً في جميع صفات النمو الخضري للفساتل المنزرعة وبمستويات معنوية مختلفة عند مقارنتها بالمعاملة العيارية في كلا الموسمين. أوضحت النتائج أيضاً أن محلول المنشط الحيوي (EM) بمفرده أعطى متوسطات نمو قريبة من تلك التي حققتها معاملة التسميد بالسماد الكيماوي الكامل بمعدل 5 جم/ كيس مع بعض الاستثناءات القليلة بكلا الموسمين. كذلك، فإن زيادة معدل إضافة السماد الكيماوي الكامل إلى 7.5 جم/ كيس أو أكثر لم تحدث أية زيادة إضافية ملحوظة في معدل النمو الخضري، بينما أدى الجمع بين المعاملة بال-EM والسماد الكيماوي المتكامل بأي معدل إلى إحداث ذلك (أي إحداث زيادة واضحة بمعدلات النمو الخضري)، خاصة عند الجمع بين المعاملة بالمنشط الحيوي (EM) بمعدل 25 مللى/ لتر والتسميد بالسماد الكيماوي المتكامل بمعدل 5 جم/ كيس، حيث أعطت هذه التوليفة أعلى المتوسطات على الإطلاق مقارنة بالمعاملات الفردية والتوليفات الأخرى بمعظم الحالات بكلا الموسمين. ولقد أمكن الحصول على نتائج مشابهة فيما يتعلق بمحتوى الوريقات من عناصر النيتروجين، الفوسفور، البوتاسيوم، وكذلك من الإندولات الكلية والتمثل الضوئي (كلوروفيل أ، ب الكاروتينويدات).

وعليه، يمكن التوصية بمعاملة فساتل نخيل البلح (صنف سيوي)، عمر سنتين والمنزرعة في أكياس بلاستيك قطرها 50 سم تحت ظروف الحقل المفتوح بمحلول المنشط الحيوي EM بمعدل 25 مل/لتر + السماد الكيماوي الكامل (20 ن: 20 فو: 20 بو + عناصر صغرى) بمعدل 5 جم/ كيس كإضافة أرضية للحصول على أفضل نمو وأعلى جودة لهذه الفساتل قبل نقلها للزراعة في الحقل.

المجلة العلمية لكلية الزراعة - جامعة القاهرة - المجلد (66) العدد الرابع (أكتوبر 2015) 336-:342.